

CD68 Recombinant antibody

Cat:B35503D**Company:** HaoKeBio**Uniprot ID:**P34810**Applications:** IHC:1:500-1:2000**Organism:**Rabbit

IHC-Polymer:1:2000-1:8000

Species reactivity:Human Mouse Rat

IHC-TSA:1:2500-1:10000

Molecular Weight Calculation: 37 kDa

IF:1:100

Observed Molecular Weight: 100-140 kDa

WB:1:1000

Background:

This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms.

Synonyms:

GP110; LAMP4; SCARD1

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Cytoplasm

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.

Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 9.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Source of Reagents:

发表[中文论文]请标注:CD68(B35503D)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:CD68(B35503D) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.